

**IN THE CLAIMS:**

*This listing of the claims will replace all prior versions and listings of claims in the application:*

1. (Canceled)

2. (Canceled)

3. (Canceled)

4. (Previously Amended)

A method of extracting total triterpenoid sapogenins from bamboo comprising:

(a) selecting bamboo material from the group consisting of Phyllostachys, Bambusa and Dendrocalamus genus of Gramineae family;

(b) preparing bamboo shaving powder having a granularity from pole, branch, leaf, shoot, shoot sheath, root or a mixture of the bamboo material;

(c) extracting free triterpenoid sapogenins from the bamboo shaving powder by mixing the bamboo shaving powder with supercritical CO<sub>2</sub> fluid and an entrainer in the amount of 5 – 15 % (v/v) of CO<sub>2</sub> until the free triterpenoid sapogenins is dissolved in the CO<sub>2</sub> fluid at temperature 50 – 60 degree C and pressure 25 – 35 Mpa;

(d) separating total triterpenoid sapogenins from the CO<sub>2</sub> fluid containing free triterpenoid sapogenins by changing the temperature of the CO<sub>2</sub> fluid to 35 – 45 degree C and the pressure to 5 – 10 Mpa to gasify the CO<sub>2</sub>;

(e) collecting a composition comprising 10 - 90% total triterpenoid sapogenins, said total triterpenoid sapogenins comprising 5 - 35% friedelin and 1 - 10% lupenone.

5. (Previously Amended)

The method of claim 4 further comprising extracting free triterpenoid sapogenins from the bamboo shaving powder circularly and dynamically with recycled CO<sub>2</sub> for 2 – 5 hours.

6. (Previously Amended)

The method of claim 4, wherein said entrainer is selected from the group consisting of methanol, ethanol, acetone and a mixture of methanol, ethanol and acetone.

7. (Previously Canceled)

8. (Currently Amended)

A method of treating ~~a disease characterized as~~ hypertension, comprising administering to a subject suffering from said ~~disease~~ hypertension a therapeutically effective amount of total triterpenoid sapogenins ~~as defined in claim 1, extracted from bamboo~~, wherein total triterpenoid sapogenins is 10-90% as determined by vanillic aldehyde and perchloric acid colorimetry using friedelin as a standard, said total triterpenoid sapogenins comprising 5-35% friedelin and 1-10% lupenone as determined by GC-MS, ~~wherein~~ said therapeutically effective amount of total triterpenoid sapogenins is administered orally in medicine or food.

9. (Currently Amended)

A method of protecting skin or hair from senescence, comprising administering to a subject a therapeutically effective amount of total triterpenoid sapogenins extracted from bamboo, wherein total triterpenoid sapogenins is 10-90% as determined by vanillic aldehyde and perchloric acid colorimetry using friedelin as a standard, said total triterpenoid sapogenins comprising 5-35% friedelin and 1-10% lupenone as determined by GC-MS to intensify SOD activity or to reduce MDA level of the skin or hair , ~~as defined in claim 1~~, wherein said therapeutically effective amount of total triterpenoid sapogenins is administered externally onto the skin or hair in a daily cosmetic.

10. (Currently Amended)

A method of in vitro inhibiting ~~preventing or treating~~ growth of cancer cells or tumor cells ~~a disease characterized as carcinoma~~, comprising treating the cancer cells or tumor cells ~~administering to a subject suffering from said disease~~ a therapeutically effective amount of total triterpenoid sapogenins extracted from bamboo, wherein total triterpenoid sapogenins is 10-90% as determined by vanillic aldehyde and perchloric acid colorimetry using friedelin as a standard, said total triterpenoid sapogenins comprising 5-35% friedelin and 1-10% lupenone as determined by GC-MS, wherein the cancer cells or tumor cells are selected from the group consisting of P<sub>388</sub> mouse leukemia cells, A<sub>549</sub> human lung adenocarcinoma cells, A<sub>375</sub> human melanoma cells, L<sub>929</sub> mouse lung epithelial tumor cells, Hela human cervical tumor cells and THP-1 human macrophage tumor cells ~~as defined in claim 1, wherein said therapeutically effective amount of total triterpenoid sapogenins is administered orally in medicine or food.~~

11. (Previously Amended)

The method according to claim 10, wherein the total triterpenoid sapogenins is pentacyclic triterpenoid sapogenins.

12. (Canceled)

13. (Canceled)

14. (Previously Amended)

The method of claim 4, wherein said granularity is 10 – 20 meshes.